

Checklist of Preventative Services

Screening tests	Coverage interval	Criteria	Criteria Met	Scheduled
Abdominal aortic aneurysm screening	One time	-FH of AAA -Men age 65-75 & smoked at least 100 cigarettes in lifetime		
Bone mass measurements	Every 24 months	-Women with Estrogen deficiency & at risk for Osteoporosis -XRay with Osteopenia, Osteoporosis or vertebral fracture -Taking Prednisone or steroid type drug -Diagnosis with primary Hyperparathyroidism -Monitoring for Osteoporosis drug therapy		
Cervical & vaginal cancer screening	-Every 24 months -12 months if at high risk	-All women		
Colorectal cancer screening -Barium enema -Colonoscopy -FOB test -Flex Sigmoidoscopy	Every -48 months -120 months -12 months -48 months	-Age 50 and older		
Mammogram screening	Once every 12 months	-Women 40 or older -Women between 35-39 can get one baseline mammogram		
Prostate cancer screening -DRE -PSA	Once every 12 months	All men over 50.		
Cardiovascular disease screening -Lipid panel	Every 5 years	-All people		
Depression screening	-one per year	-All people		
Diabetes screening	-Every year	-Any of these risk factors-HTN, Dyslipidemia, Obesity, Hyperglycemia		

		or -2 or more of these apply-Age 65 or older, overweight, FH of DM, Hx of GDM		
HIV screening	-Every year	-Increased risk -Ask for the test -Pregnant		
Obesity screening and counseling		-BMI 30 or more		
STI screening and counseling	-Every 12 months	-Pregnant -High risk for STI		
Glaucoma	-Every 12 months	-People with DM, FH of Glaucoma, African American 50 or older, Hispanic 65 or older		
Counseling services	Coverage interval	Criteria	Criteria Met	Scheduled
Nutrition therapy services		-Have DM, kidney disease, kidney transplant in last 3 years or referred by PCP.		
Alcohol misuse and screening and counseling	Every year			
Cardiovascular disease(Behavioral therapy)	One per year	-All people		
Diabetes self management training	Once with follow up every year	-At risk of complications from Diabetes		
Tobacco use cessation counseling	8 face to face visits in 12 month period	-All people who use Tobacco		
Shots	Coverage interval	Criteria	Criteria Met	Scheduled
Influenza vaccine	One per flu season	All people		
Hepatitis B vaccine		-High to medium risk for Hepatitis B-hemophilia, ESRD, DM or conditions that lower resistance to infection.		
Pneumococcal vaccine		-All people		